

Annual Symposium of the Hellenic Nuclear Physics Society

Τόμ. 28 (2021)

HNPS2021



Shell model foundations of the proxy-SU(3) symmetry: Nucleon pairs creating nuclear deformation

Dennis Bonatsos, Andriana Martinou, Ioannis E. Assimakis, Spyridon K. Peroulis, Smaragda Sarantopoulou, Nikolay Minkov

doi: [10.12681/hnps.3538](https://doi.org/10.12681/hnps.3538)

Copyright © 2022, Dennis Bonatsos, Andriana Martinou, Ioannis E. Assimakis, Spyridon K. Peroulis, Smaragda Sarantopoulou, Nikolay Minkov



Άδεια χρήσης [Creative Commons Attribution-NonCommercial-NoDerivatives 4.0](https://creativecommons.org/licenses/by-nc-nd/4.0/).

Βιβλιογραφική αναφορά:

Bonatsos, D., Martinou, A., Assimakis, I. E., Peroulis, S. K., Sarantopoulou, S., & Minkov, N. (2022). Shell model foundations of the proxy-SU(3) symmetry: Nucleon pairs creating nuclear deformation. *Annual Symposium of the Hellenic Nuclear Physics Society*, 28, 9–13. <https://doi.org/10.12681/hnps.3538>

Shell model foundations of the proxy-SU(3) symmetry: Nucleon pairs creating nuclear deformation

Dennis Bonatsos^{1,*}, Andriana Martinou¹, I.E. Assimakis¹, S.K. Peroulis¹, S. Sarantopoulou¹, N. Minkov²

¹*Institute of Nuclear and Particle Physics, National Centre for Scientific Research
“Demokritos”, Aghia Paraskevi, GR-15310 Attiki, Greece*

²*Institute of Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences, 72 Tzarigrad Road,
1784 Sofia, Bulgaria*

Abstract Proxy-SU(3) symmetry is an approximation scheme extending the Elliott SU(3) algebra of the sd shell to heavier shells, in order to make possible the application of the symmetry properties in cutting down the size of the required calculations. When introduced in 2017, the approximation had been justified by calculations carried out within the Nilsson model, an elementary shell model based on a 3-dimensional harmonic oscillator with cylindrical symmetry, applicable to deformed nuclei. Recently our group managed to map the cartesian basis of the Elliott SU(3) model onto the spherical shell model basis, fully clarifying the approximations used within the proxy-SU(3) scheme and paving the way for using the proxy-SU(3) approximation in shell model calculations for heavy nuclei. As a by-product, the relation of the 0[110] Nilsson pairs used in proxy-SU(3) to the earlier used de Shalit-Goldhaber pairs and Federman-Pittel pairs has been clarified. The connection between the proxy-SU(3) scheme and the spherical shell model has also been worked out in the original framework of the Nilsson model, with identical results.

Keywords Elliott SU(3), proxy-SU(3), shell model, de Shalit-Goldhaber pairs, Federman-Pittel pairs

BASES OF MAIN NUCLEAR STRUCTURE MODELS

The fundamental underlying model of nuclear structure is the shell model, introduced [1-4] in 1949, which is based on a three-dimensional (3D) isotropic harmonic oscillator (HO), with a spin-orbit term added to it. The states are labeled by the number of oscillator quanta n , the orbital angular momentum l , the total angular momentum j , and its z -projection m_j , with the notation $|n\ l\ j\ m_j\rangle$ being used. The shell model was considered adequate for describing near-spherical nuclei, with few valence nucleons outside closed shells.

In 1952 the collective model of Bohr and Mottelson has been introduced [5,6], in order to accommodate departure from the spherical shape and from axial symmetry, described by the collective variables β and γ respectively.

In 1955, Nilsson [7,8] introduced a modified version of the shell model, allowing for axial nuclear deformation to be included, based on a 3D anisotropic HO with cylindrical symmetry [9-14]. In the Nilsson model the states are labeled by the total number of oscillator quanta N , the number n_z of quanta along the z -axis, and the projections along the z -axis of the orbital angular momentum and the total

* Corresponding author: bonat@inp.demokritos.gr

angular momentum, represented by Λ and K respectively. Thus the Nilsson states are denoted by [7,8] $K [N n_z \Lambda]$.

In 1958 Elliott proved that deformation within the nuclear sd shell can be described in terms of the SU(3) algebra [15-19]. In contemporary language, the sd shell possesses the U(6) symmetry, while deformation within it can be described in terms of the SU(3) subalgebra of U(6). In the Elliott model the cartesian basis of a 3D isotropic HO is used, $[n_z n_x n_y m_s]$, in which the number of quanta along the z, x, y directions and the z-projection of the spin appear.

In 1972 the group theoretical structure of the Bohr--Mottelson model, having an overall U(5) symmetry possessing an O(5) subalgebra, has been understood [20]. The seniority quantum number τ , related to the irreducible representations (irreps) of O(5), plays a major role in labeling the states.

In 1975 the Interacting Boson Model [21-24] has been introduced, which also has an overall U(6) symmetry built by s-bosons of zero angular momentum and d-bosons of angular momentum two, possessing three limiting symmetries, U(5) for vibrational nuclei, which is equivalent to the Bohr-Mottelson collective model, O(6) for γ -unstable nuclei, and SU(3) for deformed nuclei.

A comprehensive review of using the SU(3) symmetry in nuclear structure has been given recently by Kota [25].

Beyond the sd nuclear shell the SU(3) symmetry of the 3D isotropic HO is known to be broken by the spin-orbit force, which within each HO shell pushes the orbitals possessing the highest angular momentum j to the shell below. As a consequence, each shell consists by the orbitals left back after this removal, called the normal parity orbitals, plus the orbitals invading from the shell above, having the opposite parity and called the intruder orbitals.

In 2017 the proxy-SU(3) symmetry has been introduced [26-28], in which the intruder orbitals in each shell (except the one with the highest projection of the total angular momentum) are replaced by the orbitals which have deserted this shell by sinking into the shell below. As a result of this replacement, each shell regains the relevant U(n) symmetry having a SU(3) subalgebra, with only one orbital (which can accommodate two particles) remaining estranged. However, this orbital is the one lying highest in energy within the shell, thus it should be empty for most of the nuclei living in this shell. Therefore, it is expected that its influence on the structure of most nuclei living in the shell should be minimal.

The proxy-SU(3) scheme has been initially justified as a good approximation through calculations [26] carried out within the Nilsson model [7,8]. Within the proxy-SU(3) scheme the importance of the highest weight irreducible representations of SU(3) has been demonstrated [29] and used [27,28] for the successful prediction of the prolate to oblate shape transition at $N=114$, the dominance of prolate over oblate shapes in the ground states of even-even nuclei, and the prediction of specific islands on the nuclear chart in which shape coexistence can appear [30]. In the present work we are going to discuss its justification through its connection to the shell model. However, before doing so, it is instructive to discuss the nature of nucleon pairs related to the development of nuclear deformation.

NUCLEON PAIRS FAVORING DEFORMATION

As early as 1953 it has been observed by deShalit and Goldhaber [31] in their studies of β transition probabilities that within the proton--neutron pairs of orbitals $(1p_{3/2}, 1d_{5/2})$, $(1d_{5/2}, 1f_{7/2})$, $(1f_{7/2}, 1g_{9/2})$, $(1g_{9/2}, 1h_{11/2})$, $(1h_{11/2}, 1i_{13/2})$ the nucleons of one kind (protons, for example) have a stabilizing effect on pairs of nucleons of the other kind (neutrons in the example), thus favoring the

development of nuclear deformation. In the standard shell model notation $|n\ l\ j\ m_j\rangle$, the orbitals forming a pair differ by $|\Delta n\ \Delta l\ \Delta j\ \Delta m_j\rangle = |0\ 1\ 1\ 0\rangle$.

A major step forward in our understanding of effective interactions and coupling schemes in nuclei has been taken in 1962 by Talmi [32] through the introduction of seniority [32-35], representing the number of nucleon pairs coupled to non-zero angular momentum, which explained the linear dependence of neutron separation energies on the mass number within series of isotopes.

In 1977 Federman and Pittel [36-38] realized that when adding valence protons and valence neutrons to a nucleus, the proton--neutron pairs (1d5/2, 1d3/2), (1g9/2, 1g7/2), (1h11/2, 1h9/2), and (1i13/2, 1i11/2) are responsible for the onset of deformation, while deformation is then established by the proton--neutron pairs (1d5/2, 1f7/2), (1g9/2, 1h11/2), (1h11/2, 1i13/2), and (1i13/2, 1j15/2). These sets correspond to $|\Delta n\ \Delta l\ \Delta j\ \Delta m_j\rangle = |0\ 0\ 1\ 0\rangle$ and $|0\ 1\ 1\ 0\rangle$ respectively in the shell model notation, the latter set coinciding with the de Shalit--Goldhaber pairs.

The decisive role played by proton-neutron pairs has been demonstrated in 1985 through the introduction of the $N_p N_n$ scheme [39,40] and the P-factor, $P = N_p N_n / (N_p + N_n)$ [41,42], by showing the systematic dependence of several observables on the competition between the quadrupole deformation, "measured" by the quadrupole-quadrupole interaction through $N_p N_n$, and the pairing interaction, "measured" through $N_p + N_n$, where N_p (N_n) is the number of valence protons (neutrons).

In 1995 the quasi-SU(3) symmetry [43,44] has been introduced, based on the proton--neutron pairs (1g9/2, 2d5/2), (1h11/2, 2f7/2), (1i13/2, 2g9/2), expressed as $|\Delta n\ \Delta l\ \Delta j\ \Delta m_j\rangle = |1\ 2\ 2\ 0\rangle$ in the shell model notation, which lead to enhanced quadrupole collectivity [45].

Following detailed studies of double differences of binding energies [46-50], in 2010 it has been realized [51] that proton-neutron pairs differing in the Nilsson notation [7,8] $K [N\ n_z\ \Lambda]$ by $\Delta K [\Delta N\ \Delta n_z\ \Delta \Lambda] = 0 [110]$, play a major role in the development of nuclear deformation, due to their large spatial overlaps [52]. These pairs correspond to the replacements made within the proxy-SU(3) scheme [26-28]. No relation to the pairs mentioned in the previous paragraphs had been realized at that time.

CONNECTING THE CARTESIAN ELLIOTT BASIS TO THE SPHERICAL SHELL MODEL BASIS

The cartesian basis of Elliott [15-19] can be transformed to the spherical basis $[n\ l\ m_l\ m_s]$ in l-s coupling through a unitary transformation [53-55]

$$[n_z\ n_x\ n_y\ m_s] = R [n\ l\ m_l\ m_s],$$

the details of which can be found in Ref. [56]. Using Clebsch-Gordan coefficients the spherical basis can be rewritten in j-j coupling as

$$[n\ l\ m_l\ m_s] = C [n\ l\ j\ m_j],$$

in which the total angular momentum j and its z-projection appear. Combining these two transformations one obtains

$$[n_z\ n_x\ n_y\ m_s] = R C [n\ l\ j\ m_j],$$

i.e., the connection between the cartesian Elliott basis and the shell model basis in j-j coupling. Details of the calculations and tables for all relevant shells can be found in Ref. [56].

Using the above transformation one sees that the Nilsson 0[110] replacements made within the proxy-SU(3) scheme are "translated" into $|0\ 1\ 1\ 0\rangle$ replacements within the spherical shell model basis.

The resulting correspondence between original shell model orbitals and proxy-SU(3) orbitals is summarized in Table 7 of Ref. [56]. This correspondence proves that the 0[110] Nilsson pairs identified

in Ref. [51] and used within the proxy-SU(3) scheme [26-28] are identical to the de Shalit--Goldhaber pairs [31] and the Federman--Pittel pairs [36-38] within the spherical shell model basis, in which they are expressed as $|0\ 1\ 1\ 0\rangle$ pairs.

The correspondence between Nilsson pairs and shell model pairs has been corroborated by calculations [57] within the Nilsson model, in which the first justification of the proxy-SU(3) scheme has been found [26]. As one can see in Tables 1 and 3 of Ref. [58], the correspondence used in proxy-SU(3) works only for the Nilsson orbitals which possess the highest total angular momentum j within their shell, which are exactly the orbitals which are replaced within the proxy-SU(3) scheme. In further corroboration of this result, a unitary transformation connecting the orbitals being replaced within the proxy-SU(3) scheme has been found [56] within the shell model basis.

This correspondence also paves the way for taking advantage of the proxy-SU(3) symmetry in shell model calculations for heavy nuclei, in a way similar to that of the symmetry-adapted no-core shell model approach [58,59] used in light nuclei.

References

- [1] M.G. Mayer, Phys. Rev. 74, 235 (1948).
- [2] M.G. Mayer, Phys. Rev. 75, 1969 (1949).
- [3] O. Haxel et al., Phys. Rev. 75, 1766 (1949).
- [4] M.G. Mayer and J.H.D. Jensen, Elementary Theory of Nuclear Shell Structure (Wiley, New York 1955).
- [5] A. Bohr, Mat. Fys. Medd. K. Dan. Vidensk. Selsk. 26, no. 14 (1952).
- [6] A. Bohr and B. R. Mottelson, Nuclear Structure Vol. II: Nuclear Deformations (Benjamin, New York 1975).
- [7] S.G. Nilsson, Mat. Fys. Medd. K. Dan. Vidensk. Selsk. 29, no. 16 (1955).
- [8] S.G. Nilsson and I. Ragnarsson, Shapes and Shells in Nuclear Structure}. (Cambridge University Press, Cambridge 1995).
- [9] Y. Takahashi, Prog. Theor. Phys. 53, 461 (1975).
- [10] R.M. Asherova et al., Nucl. Phys. A 355, 25 (1981).
- [11] G. Rosensteel and J. P. Draayer, J. Phys. A: Math. Gen. 22, 1323 (1989).
- [12] W. Nazarewicz and J. Dobaczewski, Phys. Rev. Lett. 68, 154 (1992).
- [13] W. Nazarewicz et al., AIP Conf. Proc. 259, 30 (1992).
- [14] D. Bonatsos et al., arXiv: hep-th/9411218 (1994).
- [15] J. P. Elliott, Proc. Roy. Soc. London Ser. A 245, 128 (1958).
- [16] J. P. Elliott, Proc. Roy. Soc. London Ser. A 245, 562 (1958).
- [17] J. P. Elliott and M. Harvey, Proc. Roy. Soc. London Ser. A 272, 557 (1963).
- [18] J. P. Elliott and C. E. Wildson, Proc. Roy. Soc. London Ser. A 302, 509 (1968).
- [19] M. Harvey, in M. Baranger and E. Vogt (eds) Adv. Nucl. Phys. 1, 67 (Plenum, New York, 1968).
- [20] G.N. Afanas'ev and P.P. Raychev, Fiz. Elem. Chast. At. Yadra 3, 436 (1972). [Sov. J. Nucl. Phys. 3, 229 (1972)].
- [21] A. Arima and F. Iachello, Phys. Rev. Lett. 35, 1069 (1975).
- [22] F. Iachello and A. Arima, The Interacting Boson Model}. (Cambridge University Press, Cambridge, 1987).
- [23] F. Iachello and P. Van Isacker, The Interacting Boson-Fermion Model} (Cambridge University Press, Cambridge, 1991).
- [24] A. Frank and P. Van Isacker, Symmetry Methods in Molecules and Nuclei (S y G editores, Mexico D.F., 2005).

- [25] V.K.B. Kota, *SU(3) Symmetry in Atomic Nuclei* (Springer, Singapore, 2020).
- [26] D. Bonatsos et al., *Phys. Rev. C* 95, 064325 (2017).
- [27] D. Bonatsos et al., *Phys. Rev. C* 95, 064326 (2017).
- [28] D. Bonatsos, *Eur. Phys. J. A* 53, 148 (2017).
- [29] A. Martinou et al., *Eur. Phys. J. A* 57, 83 (2021).
- [30] A. Martinou et al., *Eur. Phys. J. A* 57, 84 (2021).
- [31] A. de Shalit and M. Goldhaber, *Phys. Rev.* 92, 1211 (1953).
- [32] I. Talmi, *Rev. Mod. Phys.* 34, 704 (1962).
- [33] I. Talmi, *Nucl. Phys. A* 172, 1 (1971).
- [34] I. Talmi, *Riv. Nuovo Cim.* 3, 85 (1973).
- [35] I. Talmi, *Simple Models of Complex Nuclei* (Harwood, Chur, 1993).
- [36] P. Federman and S. Pittel, *Phys. Lett. B* 69, 385 (1977).
- [37] P. Federman and S. Pittel, *Phys. Lett. B* 77, 29 (1978).
- [38] P. Federman and S. Pittel, *Phys. Rev. C* 20, 820 (1979).
- [39] R.F. Casten, *Phys. Rev. Lett.* 54, 1991 (1985).
- [40] R.F. Casten, *Nucl. Phys. A* 443, 1 (1985).
- [41] R.F. Casten et al., *Phys. Rev. Lett.* 58, 658 (1987).
- [42] R. F. Casten, *Nuclear Structure from a Simple Perspective* (Oxford University Press, Oxford, 2000).
- [43] A. P. Zuker et al., *Phys. Rev. C* 52, R1741 (1995).
- [44] A. P. Zuker et al., *Phys. Rev. C* 92, 024320 (2015).
- [45] K. Kaneko et al., *Phys. Rev.* 103, L021301 (2021).
- [46] R.B. Cakirli et al., *Phys. Rev. Lett.* 94, 092501 (2005); erratum *Phys. Rev. Lett.* 95, 119903 (2005).
- [47] R.B. Cakirli and R.F. Casten, *Phys. Rev. Lett.* 96, 132501 (2006).
- [48] D.S. Brenner et al., *Phys. Rev. C* 73, 034315 (2006).
- [49] M. Stoitsov et al., *Phys. Rev. Lett.* 98, 132502 (2007).
- [50] R.B. Cakirli et al., *Phys. Rev. Lett.* 102, 082501 (2009).
- [51] R. B. Cakirli et al., *Phys. Rev. C* 82, 061304 (2010).
- [52] D. Bonatsos et al., *Phys. Rev. C* 88, 054309 (2013)..
- [53] K.T.R. Davies and S.J. Krieger, *Can. J. Phys.* 69, 62 (1991).
- [54] R.R. Chasman and S. Wahlborn, *Nucl. Phys. A* 90, 401 (1967).
- [55] E. Chacon and M. de Llano, *Rev. Mex. Fis.* 12, 57 (1963).
- [56] A. Martinou et al., *Eur. Phys. J. A* 56, 239 (2020),
- [57] D. Bonatsos et al., *Eur. Phys. J. Plus* 135, 710 (2020).
- [58] K.D. Launey et al., *Int. J. Mod. Phys. E* 24, 1530005 (2015).
- [59] K.D. Launey et al., *Prog. Part. Nucl. Phys.* 95, 044312 (2017).