


HNPS Advances in Nuclear Physics

Vol 30 (2024)

HNPS2023



Shape/phase transitions and shape coexistence in even-even nuclei

Dennis Bonatsos, Andriana Martinou, Spyridon Peroulis, Theodoros Mertzimekis, Nikolay Minkov

doi: [10.12681/hnpsanp.6086](https://doi.org/10.12681/hnpsanp.6086)

Copyright © 2024, Dennis Bonatsos, Andriana Martinou, Spyridon Peroulis, Theodoros Mertzimekis, Nikolay Minkov



This work is licensed under a [Creative Commons Attribution-NonCommercial-NoDerivatives 4.0](https://creativecommons.org/licenses/by-nc-nd/4.0/).

To cite this article:

Bonatsos, D., Martinou, A., Peroulis, S., Mertzimekis, T., & Minkov, N. (2024). Shape/phase transitions and shape coexistence in even-even nuclei. *HNPS Advances in Nuclear Physics*, 30, 59–66.

<https://doi.org/10.12681/hnpsanp.6086>

Shape/phase transitions and shape coexistence in even-even nuclei

D. Bonatsos^{1,*}, A. Martinou¹, S.K. Peroulis¹, T.J. Mertzimekis², N. Minkov³

¹ *Institute of Nuclear and Particle Physics, National Centre for Scientific Research "Demokritos",
Aghia Paraskevi, GR-15310 Attiki, Greece*

² *Department of Physics, National and Kapodistrian University of Athens,
Zografou Campus, GR-15784, Athens, Greece*

³ *Institute of Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences,
72 Tzarigrad Road, 1784 Sofia, Bulgaria*

Abstract Shape/phase transitions have been observed in certain regions of the nuclear chart. Shape coexistence is also known to occur in various regions of the nuclear chart, forming islands. The interrelation between these two concepts is considered in the regions around $(N=90, Z=60)$, $(N=60, Z=40)$, $(N=40, Z=34)$, in which shape coexistence due to proton-induced neutron particle-hole excitations is related to a first-order shape/phase transition from spherical to deformed shapes.

Keywords shape/phase transition, shape coexistence

A shape/phase transition (SPT) [1] corresponds to an abrupt change of the nuclear shape. In the parameter space of the Interacting Boson Model [2], a first-order SPT appears between spherical and axially symmetric deformed shapes, ending up at a point representing a second order SPT from spherical to γ -unstable (soft to triaxial deformation) shapes [3]. In the Bohr collective model framework [4], these SPTs have been described in terms of the critical point symmetries (CPS) X(5) [5] and E(5) [6], respectively. These CPS models provide parameter-independent (up to overall scales) predictions for the spectra and $B(E2)$ transition rates at the critical point. The best experimental manifestations of the X(5) CPS have been found in the $N=90$ isotones ^{150}Nd , ^{152}Sm , and ^{154}Gd [7].

Shape coexistence (SC) [8,9] is said to occur when the ground state band (gsb) is accompanied by another $K=0$ band with similar energy but very different structure. For example, one of the bands can be spherical and the other one deformed, or both bands can be deformed, one of them having a prolate shape and the other an oblate shape.

Shape coexistence has been observed in several regions of the nuclear chart, but not all over it, as it was initially expected [8]. Martinou et al. in 2021 suggested [10,11] a dual shell mechanism in the framework of the proxy-SU(3) symmetry [12-14], which is an approximation to the nuclear shell model [15,16], predicting that SC can occur only within certain stripes on the nuclear chart, depicted in Fig. 1. These predictions were in good agreement with the schematic drawings of regions in which SC has been observed experimentally, reported in the review article by Heyde and Wood in 2011 [8], as seen in Fig. 2. They are also in agreement to the more recent collection of nuclei in which SC is expected to be seen, shown in Fig. 3, based on the review article by Bonatsos et al. in 2023 [9].

In the present article, we first try to impose some quantitative limits for the appearance of SC [17]. In Table 1, data are exhibited for all nuclei beyond $Z=N=18$ for which a $K=0$ band, as well as the transition rate $B(E2; 0_2^+ \rightarrow 2_1^+)$ are known [18]. The ratios $R_{0/2} = E(0_2^+)/E(2_1^+)$ and $B_{02} = B(E2; 0_2^+ \rightarrow 2_1^+)/B(E2; 2_1^+ \rightarrow 0_1^+)$ are also shown, along with the ratio $R_{4/2} = E(4_1^+)/E(2_1^+)$, a well known indicator of collectivity.

* Corresponding author: bonat@inp.demokritos.gr

In Fig. 4(a) the data for the ratio B_{02} are plotted vs. the data for the ratio $R_{0/2}$. A clear separation is seen into two parts, a region of high B_{02} values appearing at low $R_{0/2}$ values on the left, and a region of low B_{02} values appearing at high $R_{0/2}$ values on the right, separated by the $N=90$ isotones corresponding to the X(5) CPS mentioned above. The same behavior is seen in the predictions of various theoretical models, plotted in Fig. 4(b). From Table 1 and Fig. 4 one concludes that SC can appear when the conditions $R_{0/2} < 5.7$ and $B_{02} > 0.1$ are simultaneously fulfilled.




Figure 1. Stripes in which SC can occur according to the dual shell mechanism [10,11] of the proxy-SU(3) symmetry [12-14]. Adapted from Ref. [10].




Figure 2. Regions in which SC has been observed, as given in the review article [8], are depicted in blue and are compared to the green stripes in which SC is allowed to occur, according to the dual shell mechanism [10,11] within the proxy-SU(3) model [12-14]. Adapted from Ref. [9].




Figure 3. Nuclei exhibiting SC according to the review article [9] are shown, together with azure stripes in which SC is allowed to occur according to the dual shell mechanism [10,11] within the proxy-SU(3) model [12-14], and the orange contours corresponding to $P=5$ [21], showing the borders of the regions of well deformed nuclei, depicted in yellow. Adapted from Ref. [9].




Figure 4. (a) Experimental B_{02} ratios plotted vs. R_{02} ratios. (b) The same plot for predictions by the theoretical models $X(5)$ [5], $X(5)-\beta^{2n}$ ($n=2,4,6,8$) [19], CBS [20]. Adapted from Ref. [17].

In Fig. 5 nuclei from Table 1 with ratios $R_{4/2}=E(4_1^+)/E(2_1^+)$ below 3.05 are shown by green triangles, while nuclei with $R_{4/2}$ above 3.05 are depicted by blue triangles. SC is expected for nuclei fulfilling the condition $R_{4/2}<3.05$, which lie outside the orange contours, which correspond to values of the P-factor $P=N_p N_n / (N_p + N_n)$ [21] close to 5, where N_p (N_n) is the number of valence protons (neutrons) in a given nucleus, counted from the closest closed shell. Deformed nuclei with $R_{4/2}>3.05$ lie in the yellow regions inside the orange contours and are not expected to exhibit SC. What is interesting, is

that the $N=90$ isotones being the best experimental manifestations of the $X(5)$ CPS fall on or near the orange contours, indicating that SC is indeed seen in these critical nuclei.

Table 1. Nuclei beyond $Z=N=18$ with experimentally known [18] $K=0$ bands (with energy levels given in keV) and $B(E2; 0_2^+ \rightarrow 2_1^+)$ transition rates (given in W.u.). Adapted from Ref. [17]. See text for further discussion.

nucleus	$R_{4/2}$	$E(2_1^+)$	$E(0_2^+)$	$B(E2; 2_1^+ \rightarrow 0_1^+)$	$B(E2; 0_2^+ \rightarrow 2_1^+)$	$R_{0/2}$	B_{02}
⁴⁰ Ar	1.980	1460.8	2120.9	9.0 (4)	5.3 (8)	1.452	0.589
⁴⁰ Ca		5629.4	3352.6	0.143 (+35-24)	7.5 (+30-20)	0.639	52.558
⁴² Ca	1.805	1524.7	1837.3	9.5 (4)	55. (5)	1.205	5.789
⁷⁰ Ge	2.071	1039.5	1215.6	20.8 (4)	48. (7)	1.169	2.308
⁷² Ge	2.072	834.0	691.4	23.5 (4)	89.0 (15)	0.829	3.787
⁷² Se	1.899	862.1	937.2	23.7 (17)	162. (28)	1.087	6.835
⁹⁶ Sr	2.000	814.9	1229.3	13. (8)	15.3 (16)	1.509	1.177
⁹⁸ Sr	3.006	144.2	215.6	96. (3)	62. (+7-6)	1.495	0.646
⁹⁸ Zr	1.674	1222.9	854.0	2.9 (+8-5)	145. (+40-30)	0.698	50.000
¹⁰⁰ Zr	2.656	212.5	331.1	77. (2)	67. (6)	1.558	0.870
¹⁰⁰ Mo	2.121	535.6	695.1	37.6 (9)	89. (3)	1.298	2.367
¹⁰² Mo	2.507	296.6	698.3	74. (9)	70. (30)	2.354	0.946
¹⁰⁴ Ru	2.482	358.0	988.3	57.9 (11)	25. (3)	2.761	0.432
¹¹⁰ Pd	2.463	373.8	946.7	55.5 (9)	37. (4)	2.533	0.667
¹¹² Cd	2.292	617.5	1224.3	30.31 (19)	51. (14)	1.983	1.683
¹¹⁶ Cd	2.375	513.5	1380.3	33.5 (12)	30. (6)	2.688	0.896
¹¹⁴ Sn	1.683	1299.9	1953.3	15. (3)	22. (8)	1.503	1.467
¹¹⁶ Sn	1.848	1293.6	1756.9	12.4 (4)	18. (3)	1.358	1.452
¹¹⁸ Sn	1.854	1229.7	1758.3	12.1 (15)	19. (3)	1.430	1.570
¹²⁶ Xe	2.424	388.6	1313.9	44. (4)	6.4 (12)	3.381	0.145
¹⁴⁸ Nd	2.493	301.7	916.9	57.9 (22)	3.12 (22)	3.039	0.539
¹⁵⁰ Nd	2.927	130.2	675.9	116. (3)	43.1 (23)	5.191	0.372
¹⁵² Sm	3.009	121.8	684.8	145.0 (16)	33.3 (12)	5.622	0.230
¹⁵⁴ Sm	3.255	82.0	1099.3	176. (1)	12. (3)	13.406	0.068
¹⁵² Gd	2.194	344.3	615.4	73. (+7-6)	178. (+33-53)	1.787	2.438
¹⁵⁴ Gd	3.015	123.1	680.7	157. (1)	52. (8)	5.530	0.331
¹⁵⁶ Gd	3.239	89.0	1049.5	189. (3)	8. (+4-7)	11.792	0.042
¹⁵⁸ Gd	3.288	79.5	1196.2	198. (5)	1.17 (+418-13)	15.047	0.006
¹⁶⁶ Er	3.289	80.6	1460.0	217. (5)	2.7 (10)	18.114	0.012
¹⁷² Yb	3.305	78.7	1042.9	212. (2)	3.6 (10)	13.252	0.017
¹⁷⁴ Yb	3.310	76.5	1487.1	201. (7)	1.4 (+11-5)	19.439	0.007
¹⁸⁶ Os	3.165	137.2	1061.0	93.6 (21)	0.066	7.733	0.001
¹⁹² Os	2.820	205.8	956.5	62.1 (7)	0.57 (12)	4.648	0.009
¹⁹⁶ Pt	2.465	355.7	1135.3	40.60 (20)	2.8 (15)	3.192	0.069
X(5)	2.904					5.649	0.624

In corroboration of the above findings, data are exhibited in Table 2 for all nuclei beyond $Z=N=18$ for which a $K=0$ band, as well as the transition rate $B(E2; 0_2^+ \rightarrow 2_1^+)$ are known [18], but no levels beyond 0_2^+ are known for the $K=0$ band [18]. Nuclei from Table 2 with ratios $R_{4/2}=E(4_1^+)/E(2_1^+)$ below 3.05 are added in Fig. 5 as green circles, while nuclei with $R_{4/2}$ above 3.05 are depicted by blue circles.

It should be noticed that from Fig. 5 are absent the neutron-deficient Hg, Pb, and Po isotopes, which form the first region in which SC [8,9] has been observed, the reason being that no data for their $B(E2; 0_2^+ \rightarrow 2_1^+)$ rates are shown in Ref. [18]. Existing data for these nuclei have been collected in Table 3, while these nuclei are shown as purple diamonds in Fig. 5 for completeness.

Table 2. Nuclei beyond $Z=N=18$ with experimentally known $B(E2; 0_2^+ \rightarrow 2_1^+)$ transition rates (given in W.u.) [18], but with no other levels of a $K=0$ band based on the 0_2^+ known. Energy levels are given in keV. Adapted from Ref. [17]. See text for further discussion.

nucleus	$R_{4/2}$	$E(2_1^+)$	$E(0_2^+)$	$B(E2; 2_1^+ \rightarrow 0_1^+)$	$B(E2; 0_2^+ \rightarrow 2_1^+)$	$R_{0/2}$	B_{02}
³⁸ Ar		2167.5	3376.9	3.40 (16)	1.26 (8)	1.558	0.371
⁴⁴ Ca	1.973	1157.0	1883.5	10.9 (6)	22. (7)	1.628	2.018
⁴⁸ Ca	1.175	3831.7	4283.3	1.84 (+17-14)	10.1 (5)	1.118	5.489
⁴⁶ Ti	2.260	889.3	2611.0	19.5 (6)	50. (14)	2.936	2.564
⁴⁸ Ti	2.344	983.5	2997.2	13.2 (+13-11)	20.6 (+44-32)	3.047	1.561
⁵⁰ Ti	1.722	1553.8	3868.3	5.46 (19)	1.6 (+14-5)	2.490	0.293
⁵⁴ Cr	2.185	834.9	2829.6	14.4 (6)	10. (+3-4)	3.389	0.694
⁵⁶ Fe	2.462	846.8	2941.5	16.8 (7)	2.4 (+7-12)	3.474	0.143
⁵⁸ Ni	1.691	1454.2	2942.6	10.0 (4)	0.00040 (6)	2.023	4 10 ⁻⁵
⁶⁴ Ni	1.940	1345.8	2867.3	7.76 (26)	3.15 (+23-21)	2.131	0.406
⁶⁴ Zn	2.326	991.6	1910.3	20.0 (5)	0.057 (3)	1.927	0.003
⁶⁸ Zn	2.244	1077.4	1655.9	14.69 (19)	5.5 (10)	1.537	0.374
⁷⁰ Zn	2.019	884.9	1070.8	16.7 (10)	37.3 (19)	1.210	2.234
⁷⁴ Ge	2.457	595.9	1482.8	33.0 (4)	9. (+9-6)	2.489	0.273
⁷⁴ Se	2.148	634.7	853.8	42.0 (6)	77. (7)	1.345	1.833
⁷⁶ Se	2.380	559.1	1122.3	44. (1)	47. (22)	2.007	1.068
⁷⁸ Se	2.449	613.7	1498.6	33.5 (8)	1.17 (21)	2.442	0.035
⁸⁰ Se	2.554	663.3	1478.8	24.7 (6)	6.9 (11)	2.220	0.279
⁸² Se	2.650	654.8	1410.3	17.3 (10)	3.62	2.154	0.209
⁷⁴ Kr	2.225	455.6	509.0	67. (1)	60. (17)	1.117	0.896
⁷⁸ Kr	2.460	455.0	1017.2	67.9 (22)	47. (4)	2.235	0.692
⁸² Kr	2.344	776.5	1487.6	21.3 (9)	15. (5)	1.916	0.704
⁸⁸ Sr		1836.1	3156.2	7.6 (4)	4.0 (+15-14)	1.719	0.526
⁹⁰ Zr	1.407	2186.3	1760.7	5.38 (13)	26. (50)	0.805	4.833
⁹² Zr	1.600	934.5	1382.8	6.4 (6)	14.4 (5)	1.480	2.250
⁹⁴ Zr	1.600	918.8	1300.2	4.9 (3)	9.4 (4)	1.415	1.918
⁹⁶ Mo	2.092	778.2	1148.1	20.7 (4)	51. (7)	1.475	2.464
⁹⁸ Mo	1.918	787.4	734.8	20.1 (4)	48.5 (+50-125)	0.933	2.413
⁹⁶ Ru	1.823	832.6	2148.8	18.4 (4)	12 (+5-12)	2.581	0.652
⁹⁸ Ru	2.142	652.4	1322.1	29.8 (10)	42. (+12-11)	2.026	1.409
¹⁰⁰ Ru	2.273	539.5	1130.3	35.7 (3)	35. (5)	2.095	0.980
¹⁰² Ru	2.329	475.1	943.7	44.6 (7)	35. (6)	1.986	0.785
¹⁰⁴ Pd	2.381	551.8	1333.6	36.9 (19)	13.2 (13)	2.417	0.358
¹⁰⁶ Pd	2.402	511.9	1133.8	44.3 (15)	35. (8)	2.215	0.790
¹⁰⁸ Pd	2.416	433.9	1052.8	50.4 (15)	52. (5)	2.426	1.032
¹¹⁴ Cd	2.299	558.5	1285.6	31.1 (19)	27.4 (17)	2.032	0.881
¹¹⁸ Cd	2.388	487.8	1134.5	33. (3)	5.3 (8)	2.636	0.161
¹²⁰ Sn	1.873	1171.3	1875.1	11.41 (22)	12.6 (17)	1.601	1.104
¹²⁴ Te	2.072	602.7	1657.3	31.1 (5)	20. (4)	2.750	0.643
¹²⁶ Te	2.043	666.4	1873.4	25.1 (5)	8.8 (+8-11)	2.811	0.351
¹⁴⁴ Nd	1.887	696.6	2084.7	25.9 (5)	19. (12)	2.993	0.734
¹⁵⁰ Sm	2.316	334.0	740.5	57.1 (13)	53. (5)	2.217	0.928
¹⁹⁴ Pt	2.470	328.5	1267.2	49.5 (20)	0.63 (+20-13)	3.858	0.013
¹⁹⁸ Pt	2.419	407.2	914.5	31.81 (22)	26. (7)	2.246	0.817

Table 3. Hg, Pb, and Po isotopes known to exhibit SC, for which the 0_2^+ state is experimentally known [18]. Energies are given in keV, while transition rates are given in W.u. Adapted from Ref. [17]. See text for further discussion.

nucleus	$R_{4/2}$	$E(2_1^+)$	$E(0_2^+)$	$B(E2; 2_1^+ \rightarrow 0_1^+)$	$R_{0/2}$
^{180}Hg	1.623	434.2	419.8	49. (9)	0.967
^{182}Hg	3.198	351.7	328.0	55. (3)	0.933
^{184}Hg	2.962	366.8	375.1	62. (15)	1.023
^{186}Hg	2.665	405.3	523.0	71.3 (13)	1.290
^{188}Hg	2.434	412.8	824.5	54. (9)	1.997
^{190}Hg	2.502	416.3	1278.6	45. (3)	3.071
^{184}Pb			570.0		
^{186}Pb			530.0		
^{188}Pb	1.470	723.6	591.0	7. (3)	0.817
^{190}Pb	1.588	773.9	658.0		0.850
^{192}Pb	1.588	853.6	768.8		0.901
^{194}Pb	1.596	965.1	930.7		0.964
^{196}Po	1.924	463.1	558.0		1.205
^{198}Po	1.915	604.9	816.0		1.349
^{200}Po	1.918	665.9	1136.5		1.707




Figure 5. Nuclei with $R_{4/2} < 3.05$ in which SC is expected are shown by green symbols, while nuclei with $R_{4/2} > 3.05$, for which no SC is expected, are indicated by blue symbols. Hg, Pb, and Po isotopes known to exhibit SC are shown by purple diamonds. Adapted from Ref. [17]. See text for further discussion.

In Fig. 5, nuclei in regions lighter than $N=90$ also fall on or near the orange contours. In order to examine if they do exhibit critical behavior, we need a relevant order parameter. It turns out [17,22] that the ratio $R_{2/0}=1/R_{0/2}$ is an appropriate order parameter. Indeed, using the standard IBM Hamiltonian in the consistent-Q formalism [23] with the parameter ζ playing the role of the control parameter (with

$\zeta=0$ corresponding to spherical shapes and $\zeta=1$ corresponding to deformed cases), we see in Fig. 6(a) that the ratio $R_{2/0}$ uncovers the critical point of the transition from spherical to deformed shapes, in the same way the ratio $R_{4/2}$ does, as seen in Fig. 6(b).




Figure 6. Energy ratios $R_{2/0}$ (a) and $R_{4/2}$ (b) plotted vs. the control parameter ζ , as obtained from IBM calculations using the consistent- Q formalism Hamiltonian [23]. Adapted from Ref. [17].




Figure 7. Experimental energy ratios $R_{2/0}$ in the Nd-Er (a), Sr-Pd (b), and Ni-Se (c) regions, plotted vs. the neutron number N , which serves as the control parameter. Adapted from Ref. [17].

In Fig. 7 the data for the ratio $R_{2/0}$ are shown in three different regions, with the neutron number N serving as the control parameter. The critical behavior seen in Fig. 7(a) for the $N=90$ isotones in the Nd-Er region, is also observed in Fig. 7(b) for the $N=60$ isotones in the Sr-Pd region, as well as in Fig. 7(c)

for the $N=40$ isotones in the Ni-Se region, in which SC due to proton-induced particle-hole excitations is known to occur [24,25].

In conclusion, a close connection between SPT and SC is proved to exist in the three regions ($N=90, Z=60$), ($N=60, Z=40$), and ($N=40, Z=34$). The close connection between SPTs and SC in the last two regions has been studied in detail by Heyde [26] and Garcia-Ramos [27,28].

Acknowledgments

Support by the Bulgarian National Science Fund (BNSF) under Contract No. KP-06-N48/1 is gratefully acknowledged.

References

- [1] P. Cejnar et al., *Rev. Mod. Phys.* 82, 2155 (2010); doi: 10.1103/RevModPhys.82.2155
- [2] F. Iachello and A. Arima, *The Interacting Boson Model* (Cambridge U. Press, Cambridge, 1987); doi: 10.1017/CBO9780511895517
- [3] D.H. Feng et al., *Phys. Rev. C* 23, 1254 (1981); doi: 10.1103/PhysRevC.23.1254
- [4] A. Bohr, *Dan. Mat. Fys. Medd.* 26, no. 14 (1952)
- [5] F. Iachello, *Phys. Rev. Lett.* 87, 052502 (2001); doi: 10.1103/PhysRevLett.87.052502
- [6] F. Iachello, *Phys. Rev. Lett.* 85, 3580 (2000); doi: 10.1103/PhysRevLett.85.3580
- [7] R.F. Casten and E. A. McCutchan, *J. Phys. G* 34, R285 (2007); doi: 10.1088/0954-3899/34/7/R01
- [8] K. Heyde and J.L. Wood, *Rev. Mod. Phys.* 83, 1467 (2011); doi: 10.1103/RevModPhys.83.1467
- [9] D. Bonatsos et al., *Atoms* 11, 117 (2023); doi: 10.3390/atoms11090117
- [10] A. Martinou et al., *Eur. Phys. J. A* 57, 84 (2021); doi: 10.1140/epja/s10050-021-00396-w
- [11] A. Martinou et al., *Symmetry* 15, 29 (2023); doi: 10.3390/sym15010029
- [12] D. Bonatsos et al., *Phys. Rev. C* 95, 064325 (2017); doi: 10.1103/PhysRevC.95.064325
- [13] D. Bonatsos et al., *Phys. Rev. C* 95, 064326 (2017); doi: 10.1103/PhysRevC.95.064326
- [14] D. Bonatsos et al., *Symmetry* 15, 169 (2023); doi: 10.3390/sym15010169
- [15] A. Martinou et al., *Eur. Phys. J. A* 56, 239 (2020); doi: 10.1140/epja/s10050-020-00239-0
- [16] D. Bonatsos et al., *Eur. Phys. J. Plus* 135, 710 (2020); doi: 10.1140/epjp/s13360-020-00749-2
- [17] D. Bonatsos et al., *J. Phys. G* 50, 075105 (2023); doi: 10.1088/1361-6471/acd70b
- [18] ENSDF database; url: <https://www.nndc.bnl.gov/ensdf>
- [19] D. Bonatsos et al., *Phys. Rev. C* 69, 014302 (2004); doi: 10.1103/PhysRevC.69.014302
- [20] N. Pietralla and O.M. Gorbachenko, *Phys. Rev. C* 70, 011304(R) (2004); doi:10.1103/PhysRevC.70.011304
- [21] R.F. Casten et al., *Phys. Rev. Lett.* 58, 658 (1987); doi: 10.1103/PhysRevLett.58.658
- [22] D. Bonatsos et al., *Phys. Rev. Lett.* 100, 142501 (2008); doi: 10.1103/PhysRevLett.100.142501
- [23] D.D. Warner and R.F. Casten, *Phys. Rev. Lett.* 48, 1385 (1985); doi: 10.1103/PhysRevLett.48.1385
- [24] D. Bonatsos et al., *Phys. Lett. B* 829, 137099 (2022); doi: 10.1016/j.physletb.2022.137099
- [25] D. Bonatsos et al., *Phys. Rev. C* 106, 044323 (2022); doi: 10.1103/PhysRevC.106.044323
- [26] K. Heyde et al., *Phys. Rev. C* 69, 054304 (2004); doi: 10.1103/PhysRevC.69.054304
- [27] J.-E. Garcia-Ramos and K. Heyde, *EPJ Web of Conf.* 178, 05005 (2018); doi: 10.1051/epjconf/201817805005
- [28] J.-E. Garcia-Ramos and K. Heyde, *Phys. Rev. C* 102, 054333 (2020); doi: 10.1103/PhysRevC.102.054333